

Efficacy of the bioagents *Bacillus* isolates and *Trichoderma* spp. in the control of wilt/root-rot disease in chickpea

Wilt/root-rot
disease in
chickpea

303

Hanan Ibrahim Mudawi

*The Environmental, Natural Resources and Desertification Research Institute,
National Centre for Research, Ministry of High Education,
Khartoum, Sudan, and*

Mohamed Osman Idris

*Department of Plant Protection, College of Agriculture,
Khartoum University, Khartoum, Sudan*

Abstract

Purpose – The purpose of this paper is to investigate the ability of four microorganisms to reduce disease infection of root lesion nematode (*Pratylenchus* spp.) and *Fusarium* spp. the causal agent of wilt/root-rot disease complex in chickpea.

Design/methodology/approach – A pot experiment was conducted for three consecutive winter seasons. A completely randomized design with five replicates was adopted. Two *Bacillus* isolates and *Trichoderma harzianum* and *T. viride* and their combinations were applied in infected soil. The effect on plant growth parameters, disease incidence and severity, root necrosis, weight of shoot and root, nematode population density and reproductive index were assessed.

Findings – The applications significantly ($p=0.05$) reduced the wilt/root-rot diseases complex. The reduction was attributed to the decline of the population density of nematodes in soil and root, and the suppression of the disease complex compared to controls, represented by reproductive index and the disease intensity index. Combinations were better than individual inoculation. The use of *Bacillus* isolate B3 and *T. harzianum* increased the number of flowers by 88.34 per cent. Reduction in the severity of root necrosis was in the range of 2.22-5.55 within a scale of 1-10. These findings indicate the significance of utilizing local bioagents for control of wilt/root-rot disease complex in chickpea plants.

Originality/value – The microorganisms used in this study are indigenous.

Keywords Biocontrol, *Bacillus* spp., *Cicer arietinum* L., *Fusarium* spp., *Pratylenchus* spp., *Trichoderma* spp.

Paper type Research paper

Introduction

Fusarium wilt and root-rot disease, caused by *Fusarium oxysporum* Schlechtend. Fr. *f. sp. ciceris* (Padwick) Matuo and K. Sato and *F. solani* is a major constraint to production wherever chickpea (*Cicer arietinum* L.) is grown (Haware, 1990). Many species of



This work has been published previously in its current or a substantially similar form and I have obtained permission to republish the Work. The previously published Work is fully attributed and referenced. I have provided evidence of permission to republish from the copyright holder © World Association for Sustainable Development (WASD).

This paper has been published in a conference proceedings publication by WASD in 2015. Please refer to copyright line.

root-lesion nematodes (*Pratylenchus* spp.) are very important plant parasites that affect chickpea production. The presence of these pathogens in chickpea field may interact synergistically causing more damage and higher level of infection than would result from either pathogen alone (Castillo *et al.*, 1998). Management of disease complexes appears to be less straight forward than one might anticipate. Besides chemicals, various workers suggested other control measures to replace highly toxic and potentially polluting chemicals used to control plant parasitic nematodes and fungi preferably with biological control agents and botanicals (Oostendrop and Sikora, 1989). Results from different studies showed that several strains of *Trichoderma* had a significant reducing effect on plant diseases caused by pathogens such as *Rhizoctonia solani*, *Sclerotium rolfsii*, *Phythium aphanidermatum*, *F. oxysporum*, *F. culmorum* and *Gaeumannomyces graminis* var. *tritici* under greenhouse and field conditions (Kucuk and Kivanc, 2003). The mechanisms suggested to be involved in bio-control by these fungi are antibiosis, lysis, competition, mycoparasitism and promotion of plant growth (Odeh, 2006). Soil treatment with *T. harzianum* resulted in greater growth, increased transpiration and reduced wilting index of *F. oxysporum* f.sp. *ciceris* (Siddiqui and Singh, 2004). A reduction in egg production in the root knot nematode, *Meloidogyne arenaria* were reported following soil treatments with *T. harzianum* (T.22) and *T. koningii* (T.8) (Haseeb *et al.*, 2005). In naturally infected field with *Pratylenchus thornei* combined seed treatments with *T. viride* and *Pseudomonas fluorescens* increased chickpea yield up to 29.50 per cent and the final nematode population was reduced by 22.30 per cent (Dwivedi and Johri, 2003). Whereas, simultaneous application of *P. fluorescens* plus *T. harzianum* and *T. viride* separately decreased *P. thornei* population in soil and root by 73 and 61 per cent, respectively (Bhatti and Kraft, 1992).

Besides fungi, plant-growth promoting rhizobacteria have also been used for biological control of soil-borne pathogens. They have the ability to improve plant growth by colonizing the root system and suppressing harmful rhizosphere microorganisms (Akhtar and Siddiqui, 2009), which provide biocontrol through several mechanisms such as the production of antibiotics, iron sequestering compounds, siderophores (Loper and Buyer, 1991; Siddiqui, 2000), extracellular hydrolytic enzymes, other secondary metabolites such as hydrogen cyanide and induced systemic resistance (Liu *et al.*, 1995). Siddiqui (2000) stated that “the use of *P. aeruginosa* and *B. subtilis* as seed dressing or as soil drench significantly suppressed root-rot/root-knot infection and nematode population densities under greenhouse and field conditions and thereby enhanced plant growth and yield in mungbean”. *Bacillus* spp. are known to reduce the wilting index in *F. udum* inoculated plants and improving plant growth (Haseeb *et al.*, 2005).

The present study demonstrates the use of *Bacillus* isolates and *Trichoderma* spp. in an attempt to improve the growth of the infected chickpea plants, reduce wilt/root-rot disease incidence and severity and to increase the yield of chickpea.

Material and methods

A pot experiment was conducted for three consecutive seasons during winter of 2006/2007, 2007/2008 and 2008/2009, to determine the ability of two isolates of *Bacillus* and two species of *Trichoderma*, to reduce disease infection of root lesion nematode (*Pratylenchus* spp.) and *Fusarium* spp.

Bacterial inocula

Bacillus isolates (B3 and B16) used in this study was isolated from chickpea (*C. arietinum* L.) rhizosphere. These isolates demonstrated the highest *in vitro* antagonism against

F. oxysporum f. sp. ciceris and *F. solani* according to a previous *in vitro* Petri plate bioassay. The bacterial isolates were maintained in tubes containing nutrient agar (NA) media in an incubator at 30°C. Culture tubes each containing 10 ml NA media were autoclaved for 30 min at 1 kg/cm² pressure. After the culture tubes were cooled, each tube was inoculated at 30°C with the two *Bacillus* isolates (B3 and B16). The culture tubes were then placed in an incubator for 48 h for the multiplication of the bacterial isolates. For mass production, 500 ml conical flasks containing 300 ml nutrient broth was autoclaved at the same pressure and time as before. After the flasks were cooled, each flask was inoculated with a loopful of the cultured bacterial isolates. The flasks were incubated at 28°C on an orbital shaker (Adolf Kuhner AG, Birsfelden, Switzerland) at 125 rpm for 48 h. Inoculum suspension was adjusted by serial diluting in Petri dishes containing NA media, so that the final colony forming unit count (CFU) was maintained at 1×10^6 CFU/ml.

Trichoderma spp. inocula

The two species of *Trichoderma* were cultured and maintained in PDA media. *T. harzianum* (Th) isolated from the infected chickpea fields at Shambat Research Station, and *T. viride* (Tv) was kindly provided by Plant Protection Directorate, Khartoum, Sudan. Active cultures were obtained by sub-culturing in 9 cm Petri dishes containing prepared PDA media supplemented with chloramphenicol (0.05 g/l) as bacteriostatic agent. The cultures were incubated at 25°C for seven to ten days in darkness. 10 ml of sterilized distilled water were added to each Petri dish, and the surface of the culture was scraped with a glass spreader to dislodge the spores. The spore suspension derived from three Petri dishes from each isolate was transferred separately to 300 ml sterilized flask. The concentration of the spores was determined with a haemocytometer so that the final fungal count was maintained at 1×10^6 spores/ml.

Raising and maintenance of test plants

Chickpea (*C. arietinum* L.) seeds cv. Shindi were chosen for this study on the basis of their known susceptibility reaction to *F. oxysporum f. sp. ciceris*, *F. solani* and *Pratylenchus* spp (Abbo, 2004). Seeds were surface sterilized with 2 per cent sodium hypochlorite for 2 min and rinsed three times in sterilized water. Earthen pots, 20 cm in diameter, were sterilized in oven at 80°C for 48 h. Each pot was filled with 2.5 kg of infected soil obtained from the wilt sick plot at Shambat Research Station, Khartoum, Sudan. The control treatment comprises two lots of pots. Five pots were filled with infected soil and the other five pots were filled with autoclaved soil at 80°C for 2 h. Ten seeds were sown per pot. The pots were kept in green house. Seeds were placed in the soil surface approximately 2 cm from the top of the pot. Each pot received 10 ml of each of *Bacillus* isolate (B3 and B16) and/ or *Trichoderma* spp. (Tv and Th) applied around the chickpea seeds and covered with soil, then immediately irrigated. Control plants received an equivalent amount of sterilized water. The treatments were as follows: B16 alone, B3 alone, Tv alone, Th alone, B16 × Tv, B16 × Th, B3 × Tv, B3 × Th, control infected soil (CI), and control sterilized soil. Two weeks after planting, seedlings were thinned to three plants per pot, selected for uniformity in growth. Plants watered daily and fertilized with nitrogen fertilizer once at the rate of 50 kg/feddan.

Disease assessment

The disease reaction was assessed by the incidence and severity of *Fusarium* spp. and Nematode symptoms on individual plants at seven to eight days intervals until the end of the experiment. The incidence of foliar symptoms (I), rated 0 or 1, was recorded

according to the number of plants infected. The severity of *Fusarium* wilt was rated on a scale from 0 to 4, according to percentage of foliage with yellowing or necrosis (0 = 0 per cent, 1 = 1-33 per cent, 2 = 34-66 per cent, 3 = 67-100 per cent, 4 = death of the plant) adopting the scale proposed by Landa *et al.* (2001) and Castillo *et al.* (2003). The incidence of foliar symptoms (*I*) and severity data (*S*) were used to calculate a disease intensity index (DII) as follows: $DII = (I \times S)/4$. Thus, DII expresses the mean value of disease intensity at any given moment as a proportion of the maximum possible amount of disease (Castillo *et al.*, 2003).

The plant growth parameters are:

- (1) plant height;
- (2) number of leaves;
- (3) number of branches; and
- (4) number of flowers.

All parameters were recorded each two weeks after seedling emergence. The fresh weight and dry weight were also determined and recorded.

The experiment was carried out in a completely randomized design with five replicates. Duncan's multiple range test, was used to test significant differences between treatments at $p \leq 0.05$ (Gomez and Gomez, 1984).

Observations

The plants were uprooted from the pots 105 days after inoculation and the root systems were rinsed gently. Each plant was cut with a knife above the base of the root-emergence zone to separate shoot and root. Shoot and root fresh weights were recorded. Root necrosis was assessed on a scale of 0-10 according to the percentage of necrotic tissue visible, in which 0 = 0 per cent, 1 = 1-10 per cent, and 10 = 91-100 per cent (Castillo *et al.*, 1998). The nematode population density in soil and roots was assessed. A 100 g subsample of well-mixed soil from each treatment was processed by Baermann tray extraction method (Southey, 1986; Hooper *et al.*, 2005). Nematode suspensions were collected in 500 ml flasks after 24 h and the numbers of nematodes counted in five aliquots of 1 ml suspension from each sample were recorded. The means of five counts were used to calculate the population of nematodes per kg of soil. To calculate the nematode number inside the roots, a 1 g subsample of roots was cut into small pieces 1 cm in length, and centrifuged in 10 ml of sterilized water for 4-5 min at 1,750 rpm. After spinning the suspension containing nematode were poured into vials and stored at 4°C for nematode examination, identification and were counted under a stereomicroscope (Siddiqui, 2000; Samac and Kindel, 2001). The numbers of nematodes present in the roots were determined per 1 g of root. The average population densities were used to calculate the reproduction index ($Rf = \text{final population density} / \text{initial population density}$) (Castillo *et al.*, 1995). For dry-weight determination, shoots were kept in envelopes at 80°C in oven for 48 h and weight was recorded in grams.


Results

Bacillus isolates and *Trichoderma* spp. used in the experiment has a great potential as biocontrol agents (BCA) for wilt-root/rot disease complex management when incorporated together rather than alone. These BCAs were successful in increasing plant growth parameters and weights, and reducing the infestation and population density of the root lesion nematode *Pratylenchus* spp. in soil and roots indicated by the DII.

Inoculation with *Bacillus* isolates and *Trichoderma* spp. and their combinations induced a significant increase ($p \leq 0.05$) in the chickpea growth parameters vs plant height, the number of branches, leaves and flowers compared to the CI plants. Inoculation with *T. harzianum* increased plant height by 53.32 per cent over the others inoculations; however the greatest increase in number of leaves and flowers were recorded when *Bacillus* isolate B16 was incorporated with *T. viride* (47.42 per cent, 88.83 per cent, respectively). Inoculating with *T. viride* alone increased the number of branches by 51.32 per cent (shown in Figure 1).

On average plant inoculation with *T. harzianum* produced a pronounced reduction effect on almost all the tested parameters effecting nematode progress. It reduced the number of nematodes in soil and roots (30.21 and 75.63 per cent, respectively) with a DII of 0.52 compared to 3.29 in CI plants (Table I). When *Bacillus* isolate B3 was combined with *T. harzianum*, it reduced the severity of root necrosis by 2.74 and the reproductive factor by 1.19 compared with CI plants (8.16 and 5.94, respectively) (Table II). No significant reduction levels were recorded between all the tested antagonisms in the DII. However, the applied combination of *Bacillus* isolate B16 and *T. viride* gave the least reduction level of 0.44 (Table II).

Chickpea fresh root and dry shoot weights were highly affected when inoculated with the mixed application with no significant difference ($p \leq 0.05$) between applications. The application of *T. harzianum* and *Bacillus* isolate B16 was the most



Notes: Error bares represent standard error for means. Each point is the mean of five replicates for three seasons (2006/2007-2007/2008-2008/2009). (a) Plant height; (b) number of branches, (c) number of leaves; (d) number of flowers per plant, compared to control sterilized (CS) and infected plants (CI)

Figure 1.
The percentage
effect of *Bacillus*
isolate B16, and B3,
Trichoderma viride
(Tv) and *T.*
harzianum (Th) with
their combinations

Table I.
Effect of *Bacillus*
isolates B16 and B3,
Trichoderma viride
(Tv) and *T.*
harzianum (Th) and
their combinations
on the number of the
nematode
Pratylenchus spp. in
soil and roots

effective application, recording a high increase in fresh root and dry shoot weight percentages of 58.22 and 53.60 per cent, respectively (Figure 2).

The effect of the bacterial and fungal antagonisms on the disease progress of *Fusarium* wilt on chickpea plants, as illustrated in Figure 3, demonstrated that all the applied antagonisms have the ability to decrease the progress of the *Fusarium* wilt disease from day 15 after inoculation forward. However, the combined applications were more highly effective in reducing the disease progress curve than the individual ones, especially *Bacillus* isolate B16 when combined with the two *Trichoderma* spp. (Figure 3).

Discussion

In the current study, Chickpea plants (*Cicer arietinum* L.), variety Shindi was found highly susceptible to *Fusarium* wilt/root-rot diseases complex, as indicated by the high records of root necrotic severity, nematode population densities and plant growth suppression in comparison with the controls. The present results indicated that the

Treatments	Percentage number of nematodes (%)	
	In soil	In roots
B16	61.87cd	22.04b
B3	59.56c	18.72b
Tv	60.23c	19.36b
Th	69.79cd	24.36b
B16 × Tv	52.05bc	19.64b
B16 × Th	45.03bc	19.48b
B3 × Tv	40.66bc	19.49b
B3 × Th	36.77b	16.32b
CI	100.00e	100.00c
CS	0.00a	0.00a

Note: Means followed by the same letter are not significantly different at $p \leq 0.05$ according to Duncan's multiple-range test

Table II.
Effect of *Bacillus*
isolates B16 and B3,
Trichoderma viride
(Tv) and *T.*
harzianum (Th) and
their combinations
on the severity of
root necrosis, the
reproduction index
(Rf) and the disease
intensity in chickpea
plants


Treatments	Severity of root necrosis ^a	Rf % ^b	DII ^c
B16	3.88bcd	53.68c	0.56b
B3	4.44d	55.18cd	0.55b
Tv	4.31d	53.30c	0.52b
Th	4.06bcd	60.78d	0.45b
B16 × Tv	3.33bcd	43.88bc	0.44b
B16 × Th	3.20bcd	34.97ab	0.54b
B3 × Tv	4.22cd	29.00ab	0.52b
B3 × Th	2.74b	22.09a	0.45b
CI	8.16e	100.00e	3.29c
CS	0.00a	0.38a	0.00a

Notes: Means followed by the same letter are not significantly different at $p \leq 0.05$ according to Duncan's multiple-range test. ^aSeverity of root necrosis was assessed on a scale of 0-10 according to the following scale, 0 = 0 per cent, 1 = 1-10 per cent, and 10 = 91-100 per cent; ^bRf (the reproduction index) = final population density divided by initial population density; ^cDII (the disease intensity index) = incidence of foliar symptoms × severity data/4, after 90days of inoculation

combination of the BCA was more effective than either used alone in reducing disease progress and/or promoting the plant growth. As reported by Duffy *et al.* (1996), Larkin and Fravel (1998), Howell (2003) and Maketon *et al.* (2008) the use of combination of multiple antagonistic organisms may provide improved disease control over the use of single organism. They enhance the level and consistency of control by providing multiple mechanisms of action, more stable rhizosphere community, and are effective over a wider range of environmental conditions. In particular, combinations of fungi and bacteria may provide protection at different times or under various conditions, and occupy different or complementary niches. Such combinations may overcome inconsistencies in the performance of individual isolates. A single biocontrol agent may not perform well at all times in all kinds of soil environment to suppress plant pathogens (Raupach and Kloepper, 1998).

The mixed inoculation of *Bacillus* isolates and *Trichoderma* spp. enhanced plant growth expressed in an increase in plant height, number of branches, leaves, flowers, and fresh and dry matter accumulation and the suppression of the disease complex up to 100 per cent compared to controls. This is represented by severity of root necrosis and DII, better than individual application and in less time period. The promotion of chickpea growth parameters by *Bacillus* isolates and *Trichoderma* spp. may be due to their abilities to produce phytohormones, vitamins and solubilising minerals besides, their role in direct inhibition of pathogen growth (Morsy *et al.*, 2009).

The individual effect of the *Bacillus* isolates in promoting plant growth was rarely observed (Figure 1). The combined effect of *T. harzianum* with either of the *Bacillus* isolates was more effective than the inoculation with *T. viride*, which was evidenced in the enhanced growth in most of the parameters and disease reduction percentages. These results are in agreement with Abeysinghe's (2007) findings who observed that protection of bean seedlings was more pronounced in *T. harzianum* RU01 treated plants than bacterized plants with *B. subtilis* CA32 and enhanced root growth was observed only in *T. harzianum* RU01 treated plants. This suggests that biotic modifications of the mycorrhizosphere were a result of colonization with *T. harzianum* RU01.



Notes: Each column is the mean of five replicates for three seasons (2006/2007-2007/2008-2008/2009). Vertical bars represent standard errors

Figure 2.
The percentage effect of *Bacillus* isolates (B16 and B3), *Trichoderma viride* (Tv) and *Trichoderma harzianum* (Th) and their combinations on fresh root weight and dry shoot weight on chickpea susceptible plants




Figure 3.
Disease progress of
Fusarium wilt in
chickpea plants
grown in infected
soil and treated with
Bacillus isolate B16

Notes: Each drop lines represent the progress of the disease compared to control (CI). Each point is the mean of disease intensity index of five replicates for three seasons (2006/2007-2007/2008-2008/2009); each replicate consists of three potted plants. (a) *Bacillus* isolate B3; (b) *Trichoderma viride*; (c) *T. harzianum*; (d) their combinations compared to control sterilized (CS) and infected plants (CI)

The findings of Maketon *et al.* (2008) showed that neither *B. subtilis* AP-01 nor *T. harzianum* AP-001 alone could control the bacterial wilt, damping-off and frog-eye leaf spot of tobacco diseases, but when combined, their controlling capabilities increased. Monte (2001) explained this finding as a result of augmentation of levels of inhibitory compounds, which was not seen when either antagonism was applied alone. Hence *Trichoderma* strains strongly stimulate plants to produce their own antimicrobial compounds.

The *Trichoderma* spp. survives in the soil by using organic matter as their primary source of nutrients and nematodes as a secondary source. Thus, their behaviour in soil ecosystem to some extent mirrors that of predacious fungi that use nematodes as a food base when under nutritional stress (Sikora *et al.*, 2005). The fungi's lack of dependence on nematodes for growth and survival in soil is reflected by the fact that the DII is not correlated with nematode density (Tables I and II and Figure 3). This finding was recorded when chickpea plants were treated with *Trichoderma* spp. In this study, the inoculation of *Trichoderma* spp. singly was less effective in nematode control compared to the mixed applications. Organic matter composition and the associated biotic and abiotic factors can affect the activities of *Trichoderma*, especially in relation to the conduciveness/receptivity of the soil to the strain (Vinale *et al.*, 2008).

The mechanisms of action used by *Trichoderma* – competition, antibiosis, parasitism and systemic-induced resistance – are influenced by concentration and

availability of nutrients (carbohydrates in lignocellulosic substances, chitin, lipids, etc.) within the soil organic matter (Hoitink *et al.*, 2006).

Moreover, a high percentage of nematode population density in soil was recorded even though a minimum percentage of the average of nematode population density in soil was estimated (Table I). These were higher than what Castillo *et al.*'s (1998) estimation (1,000 nematodes/100 g) in infected chickpea plants with lesion nematodes, irrespective of fungal inoculum or nematode densities.

The high percentage of nematode population density within soils (36.77-69.79 per cent) compared to that in roots (16.32-24.36 per cent) (Table I) may be attributed to the induced resistance capacity of both antagonists (Castillo *et al.*, 1998; Hoitink *et al.*, 2006) and to the explanation of Monte (2001), although the number of nematodes in roots was lower compared to chickpea susceptible lines and cultivars used in Castillo experiments (Castillo *et al.*, 1995). According to an experiment conducted by Siddiqui *et al.* (2001) they stated that:

[...] despite reduction, final populations remained at fairly high levels, presumably because short-duration colonization results in delayed nematode hatch and ultimate penetration.

In addition, only a portion of the roots and root tips are colonized with the bacterial and fungal antagonisms and therefore a certain amount of root penetration always occurs. The combined inoculation of *T. harzianum* with *Bacillus* isolate B16 resulted in great increase in fresh root and dry shoot weight and reduced the population density of *Pratylenchus* spp. in soil and roots by 73.38 and 86.94 per cent, respectively compared to that reported by Bhatti and Kraft (1992), who stated that "simultaneous application of *P. fluorescens* plus *T. harzianum* and *T. viride* separately decreased *P. thornei* population in soil and root by 73% and 61%, respectively".

All the treatments were effective in reducing the disease, indicated by the reduced severity of root necrosis which was in the range of 2.71-4.44 in treated plants, whereas the range in the infected soil reached up to 8.3. Even the nematode reproductive index, indicated by Rf, was highly reduced by the co-application of *Bacillus* isolate B3 with *T. harzianum* with a reduction range of 22.09 per cent. This is more effective than *P. fluorescens* plus *T. viride*, when applied to control *P. thornei* in chickpea plants which resulted in a final nematode population reduction of 22.30 per cent (Dwivedi and Johri, 2003). Most rhizobacteria act against plant-parasitic nematodes by means of metabolic by-products, enzymes and toxins. The effects of those toxins include the suppression of nematode reproduction, egg hatching and juvenile survival, as well as direct killing of nematodes (Tian *et al.*, 2007). Moreover, Morsy *et al.* (2009) reported that *B. subtilis* can secrete several antifungal metabolites such as subtilin, bacitracin, bacillin and bacillomycin, which have an inhibitory effect on fungal pathogens.

The fungal and bacterial antagonists survive in the soil by using organic matter as their primary source of nutrients and nematodes as a secondary source and use nematodes as a food base when under nutritional stress and their lack of dependence on nematodes for growth and survival in soil is reflected by the fact that the severity of root necrosis is not correlated with nematode density (Sikora *et al.*, 2005).

Conclusions

According to the findings of this study, the following conclusions can be drawn:

- the use of the selected plant growth promoting rhizobacteria *Bacillus* isolates and the fungal antagonism *Trichoderma* spp. are antagonistic to both fungi and nematodes; and

- they could be developed into a valuable crop management programme to reduce the deleterious impact of soil-borne pathogenic fungi and plant-parasitic nematodes on plant growth.

Successful use of microbial antagonists will depend first upon obtaining a thorough understanding of their ecological behaviour, followed by studies of their physiology and genetics.

Future research studies should concentrate on the metabolites the microbes produce and what happen when they are inoculated in combination, since individual inoculation of these microbes gave less control of plant diseases.

References

- Abbo, A.S.H. (2004), "A study on wilt/root-rot complex in chickpea (*Cicer arietinum* L.) in Shambat area", MSc (Agriculture), thesis, Faculty of Agriculture, University of Khartoum.
- Abeysinghe, S. (2007), "Biological control of *Fusarium solani* f. sp. *phaseoli* the causal agent of root rot of bean using *Bacillus subtilis* CA32 and *Trichoderma harzianum* RU01", *Ruhuna Journal of Science*, Vol. 2 No. 1, pp. 82-88.
- Akhtar, M.S. and Siddiqui, Z.A. (2009), "Use of plant growth-promoting rhizobacteria for the biocontrol of root-rot disease complex of chickpea", *Australasian Plant Pathology*, Vol. 38 No. 2, pp. 44-50.
- Bhatti, M.A. and Kraft, J.M. (1992), "Effect of inoculum density and temperature on root rot and wilt of chickpea", *Plant Disease*, Vol. 76 No. 9, pp. 960-963.
- Castillo, P., Jiménez-Díaz, R.M., Gomez-Barcina, A. and Vovlast, N. (1995), "Parasitism of the root-lesion nematode *Pratylenchus thornei* on chickpea", *Plant Pathology*, Vol. 44 No. 4, pp. 728-733.
- Castillo, P., Mora-Rodríguez, M.P., Navas-Cortés, J.A. and Jiménez-Díaz, R.M. (1998), "Interactions of *Pratylenchus thornei* and *Fusarium oxysporum* f. sp. *ciceris* on chickpea", *Phytopathology*, Vol. 88 No. 8, pp. 828-836.
- Castillo, P., Navas-Cortés, J.A., Gomar-Tinoco, D., Di Vito, M. and Jiménez-Díaz, R.M. (2003), "Interactions between *Meloidogyne artiellia*, the cereal and legume root-knot nematode, and *Fusarium oxysporum* f. sp. *ciceris* race 5 in chickpea", *Phytopathology*, Vol. 93 No. 12, pp. 1514-1523.
- Duffy, B.K., Simon, A. and Weller, D.M. (1996), "Combination of *Trichoderma koningii* with fluorescent pseudomonads for control of take-all on wheat", *Phytopathology*, Vol. 86 No. 2, pp. 188-194.
- Dwivedi, D. and Johri, B.N. (2003), "Antifungals from fluorescent pseudomonads: biosynthesis and regulation", *Current Science*, Vol. 85 No. 12, pp. 1693-1703.
- Gomez, K.A. and Gomez, A.A. (1984), *Statistical Procedures for Agriculture Research*, 2nd ed., Wiley, New York, NY, pp. 680.
- Haseeb, A., Sharma, A. and Shukla, P.K. (2005), "Studies on the management of root-knot nematode, *Meloidogyne incognita* – wilt fungus, *Fusarium oxysporum* disease complex of green gram, *Vigna radiata* cv ML-1108", *Journal of Zhejiang University Science*, Vol. 6 No. 8, pp. 736-742.
- Howare, M.P. (1990), "Fusarium wilt and other important diseases of chickpea in the Mediterranean area", in Saxena, M.C., Cubero, J.I. and Wery, J. (Eds), *Present Status and Future Prospects of Chickpea Crop Production and Improvement in the Mediterranean Countries*, CIHEAM, Zaragoza.
- Hoitink, H.A.J., Madden, L.V. and Dorrance, A.E. (2006), "Systemic resistance induced by *Trichoderma* spp.: interactions between the host, the pathogen, the biocontrol agent, and soil organic matter quality", *Phytopathology*, Vol. 96 No. 2, pp. 186-189.

- Hooper, D.J., Hallmann, J. and Subbotin, S.A. (2005), "Methods for extraction, processing and detection of plant and soil nematodes", in Luc, M., Sikora, R.A. and Bridge, J. (Eds), *Plant Parasitic Nematodes in Subtropical and Tropical Agriculture*, 2nd ed., CAB International, Wallingford, pp. 53-86.
- Howell, C.R. (2003), "Mechanisms employed by *Trichoderma* species in the biological control of plant diseases: the history and evolution of current concepts", *Plant Disease*, Vol. 87 No. 1, pp. 4-10.
- Kucuk, C. and Kivanc, M. (2003), "Isolation of *Trichoderma* spp. And determination of their antifungal, biochemical and physiological features", *Turkish Journal of Biology*, Vol. 27 No. 4, pp. 247-253.
- Landa, B.B., Navas-Cortés, J.A., Hervás, A. and Jiménez-Díaz, R.M. (2001), "Influence of temperature and inoculum density of *Fusarium oxysporum* f. sp. *ciceris* on suppression of *Fusarium* wilt of chickpea by rhizosphere bacteria", *Phytopathology*, Vol. 91 No. 8, pp. 807-816.
- Larkin, R.P. and Fravel, D.R. (1998), "Efficacy of various fungal and bacterial biocontrol organisms for control of *Fusarium* wilt of tomato", *Plant Disease*, Vol. 82 No. 9, pp. 1022-1028.
- Liu, L., Kloepper, J.W. and Tuzun, S. (1995), "Induction of systemic resistance in cucumber against *Fusarium* wilt by plant growth promoting rhizobacteria", *Phytopathology*, Vol. 85 No. 6, pp. 695-698.
- Loper, J.E. and Buyer, J.S. (1991), "Siderophore in microbial interactions on plant surfaces", *Molecular Plant-Microbe Interactions*, Vol. 4 No. 1, pp. 5-13.
- Maketon, M., Apisitsantikul, J. and Siriraweeikul, C. (2008), "Greenhouse evaluation of *Bacillus subtilis* ap-01 and *Trichoderma harzianum* ap-100 in controlling tobacco diseases", *Brazilian Journal of Microbiology*, Vol. 39 No. 2, pp. 296-300.
- Monte, E. (2001), "Understanding *Trichoderma*: between biotechnology and microbial ecology", *International Microbiology*, Vol. 4 No. 1, pp. 1-4.
- Morsy, E.M., Abdel-Kawi, K.A. and Khalil, M.N.A. (2009), "Efficiency of *Trichoderma viride* and *Bacillus subtilis* as biocontrol agents against *Fusarium solani* on tomato plants", *Egyptian Journal of Phytopathology*, Vol. 37 No. 1, pp. 47-57.
- Odebode, A.C. (2006), "Control of postharvest pathogens of fruits by culture filtrates from antagonistic fungi", *Journal of Plant Protection Research*, Vol. 46 No. 1, pp. 1-5.
- Oostendrop, M. and Sikora, R.A. (1989), "Utilization of antagonistic rhizobacteria as seed treatment for the biological control of *Heterodera schachtii* in sugarbeet", *Revue de Nematologie*, Vol. 12 No. 1, pp. 77-83.
- Raupach, G.S. and Kloepper, J.W. (1998), "Mixtures of plant growth promoting rhizobacteria enhance biological control of multiple cucumber pathogens", *Phytopathology*, Vol. 88 No. 11, pp. 1158-1164.
- Samac, D.A. and Kindel, L.L. (2001), "Suppression of the root lesion nematode (*Pratylenchus penetrans*) in alfalfa (*Medicago sativa*) by *Streptomyces* spp.", *Plant Soil*, Vol. 235 No. 1, pp. 35-44.
- Siddiqui, A.I., Ehetshamul- Haque, S. and Shaikat, S. (2001), "Use of rhizobacteria in the control of root rot-root knot disease complex of mungbean", *Journal of Phytopathology*, Vol. 149 No. 6, pp. 337-346.
- Siddiqui, M.R. (2000), *Tylenchida, Parasites of Plants and Insects*, 2nd ed., CAB International, Wallingford, pp. 30-34.
- Siddiqui, Z.A. and Singh, L.P. (2004), "Effects of soil inoculants on the growth, transpiration and wilt disease of chickpea", *Journal of Plant Diseases and Protection*, Vol. 111 No. 2, pp. 151-157.

- Sikora, R.A., Greco, N. and Veloso Silva, J.F. (2005), "Nematode parasites of food legumes", in Luc, M.R., Sikora, A. and Bridge, J. (Eds), *Plant Parasitic Nematodes in Subtropical and Tropical Agriculture*, CABI Publishing, Wallingford, pp. 259-318.
- Southey, J.F. (1986), *Laboratory Methods for Work with Plant and Soil Nematodes*, Ministry of Agriculture, Fisheries and Food, London, p. 202.
- Tian, B., Yang, J. and Zhang, K. (2007), "Bacteria used in the biological control of plant-parasitic nematodes: populations, mechanisms of action, and future prospects", *FEMS. Microbiology Ecology*, Vol. 61 No. 2, pp. 197-213.
- Vinale, F., Sivasithamparam, K., Ghisalberti, E.L., Marra, R., Woo, S.L. and Lorito, M. (2008), "Trichoderma plant-pathogen interactions", *Soil Biology and Biochemistry*, Vol. 40 No. 1, pp. 1-10.

About the authors

Dr Hanan Ibrahim Mudawi is a Researcher Professor Assistant in Bio-pesticides and Plant Pathology, currently working at the Department of Bio-pesticides and Bio-fertilizers, Environment and Natural Resources and Desertification Research Institute, National Center of Research, Ministry of High Education, Khartoum, Sudan. She was awarded a PhD in Bio-pesticides by the Sudan Academy of Sciences, and achieved her MSc (plant protection) at the University of Khartoum and BSc (Honours) in Agricultural Sciences at the University of Gazera. She has held, organized and attended many training courses in different fields and supervised undergraduate and postgraduate students. She also participated in many projects dealing with organic agriculture, a Member of some scientific societies and published some papers in peer-reviewed journals. She is a Consultant for private companies in the field of plant pathology and nematodes. Dr Hanan Ibrahim Mudawi is the corresponding author and can be contacted at: hananmodawi@gmail.com

Dr Mohamed Osman Idris is an Associate Professor in Plant Pathology currently working at the College of Agriculture, Khartoum University, Khartoum, Sudan. He has been a Lecturer on Plant Pathology since 1975. He has been awarded his PhD at the Reading University, England, MSc at the University of Khartoum and BSc (Honours) at the University of Khartoum. He is an internal and external examiner for postgraduate programmes at many universities in Khartoum. Till now he supervised studies for 55 MSc Degrees and five PhD Degrees. He has published 22 papers in national and international reputable peer-reviewed and conferences. His main research theme is fungal diseases and mycotoxins in Sudan Agricultural Products. He is the Chairman of Mycotoxins Committee Sudanese Standards and a Member of the Metrology Organization. And is currently a Consultant to many governmental and private organizations in Sudan.